



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

The following is the latest report on plague in Peru received from the Director de Salubridad:

Locality.	Cases Mar. 27.	New.	Recovered.	Died.	Remaining Apr. 3.
Lima.....	8	6	1	2	11
Chosica.....	3		2		1
Callao.....	1		1		
Trujillo.....	23	8	3	7	21
San Pedro.....	4		1		3
Paíta.....	10	11	5	5	11
Chiclayo.....	8	1?	1	?	7
Lambayeque.....	4	7		3	8
Eten.....	2	6		5	3
Santiago de Cao.....		1		1	

The figures given for Eten in last week's report were incorrect owing to an error in transmission of the report from that place.

#### PHILIPPINE ISLANDS.

##### *Report from Manila—Smallpox—Inspection of vessels.*

Chief Quarantine Officer Heiser reports, March 27:

Week ended March 23, 1907. Quarantinable disease reported for the city of Manila as follows: Smallpox, 7 cases, 0 deaths.

During the week the following vessel cleared for United States ports:

On March 20, the British steamship *Fowley*, with 51 crew, en route from New York to Cebu, was granted a supplemental bill of health.

#### FOREIGN AND INSULAR STATISTICAL REPORTS OF COUNTRIES AND CITIES—UNTABULATED.

**ALGERIA—Algiers.**—Month of March, 1907. Estimated population, 155,000. Total number of deaths, 381, including 3 from diphtheria, 1 from enteric fever, 1 from measles, 8 from smallpox, and 81 from tuberculosis.

**FRANCE—Nice.**—Month of March, 1907. Estimated population, 130,000. Total number of deaths not reported. Two deaths from enteric fever, 6 from smallpox, and 22 from tuberculosis reported.

**St. Etienne.**—Two weeks ended March 31, 1907. Estimated population, 150,000. Total number of deaths, 165, including 2 from diphtheria.

**GERMANY—Weimar.**—Month of March, 1907. Estimated population, 31,127. Total number of deaths, 60, including diphtheria 3, enteric fever 1, and 6 from tuberculosis.

**GREAT BRITAIN—England and Wales.**—The deaths registered in 76 great towns in England and Wales during the week ended April 13, 1907, correspond to an annual rate of 16.9 per 1,000 of population, which is estimated at 16,024,458.

*London.*—One thousand four hundred and ninety-five deaths were registered during the week, including measles 45, scarlet fever 9, diphtheria 8, enteric fever 5, whooping cough 60, and 18 from diarrhea. The deaths from all causes correspond to an annual rate of 16.4 per 1,000. In Greater London 2,096 deaths were registered. In the "outer ring" the deaths included 7 from diphtheria, 10 from measles, 13 from whooping cough, and 3 from scarlet fever.

*Ireland.*—The average annual death rate represented by the deaths registered during the week ended April 13, 1907, in the 21 principal town districts of Ireland was 23.5 per 1,000 of the population, which is estimated at 1,117,547. The lowest rate was recorded in Ballymena, viz, 4.8, and the highest in Kilkenny, viz, 49.1, per 1,000. In Dublin and suburbs 207 deaths were registered, including diphtheria 2, measles 7, enteric fever 1, whooping cough 7, and 31 from tuberculosis.

*Scotland.*—The deaths registered in 8 principal towns during the week ended April 13, 1907, correspond to an annual rate of 20.2 per 1,000 of the population, which is estimated at 1,812,171. The lowest rate of mortality was recorded in Paisley, viz, 14.1, and the highest in Glasgow, viz, 23.6, per 1,000. The aggregate number of deaths registered from all causes was 700, including diphtheria 5, enteric fever 4, measles 8, scarlet fever 2, typhus fever 1, and 55 from whooping cough.

*JAPAN—Formosa.*—Two weeks ended April 6, 1907. Estimated population, 3,050,004. Total number of deaths not reported. One hundred and fifty-nine deaths from plague reported.

*MALTA.*—Two weeks ended April 6, 1907. Estimated population, 205,059. Total number of deaths, 166, including 3 from diphtheria.

*NORWAY—Bergen.*—Month of March, 1907. Estimated population, 80,000. Total number of deaths, 25, including diphtheria 6, enteric fever 1, and 2 from whooping cough.

*SPAIN—Almeria.*—Month of March, 1907. Estimated population, 46,443. Total number of deaths, 118, including measles 4, smallpox 13, and 8 from tuberculosis.

*VENEZUELA.*—Reports from the entire Republic for the year ending December 31, 1905, show as follows: Estimated population, 2,323,527. Total number of deaths, 58,100, including diphtheria 97, enteric fever 1,482, measles 46, scarlet fever 72, whooping cough 582, typhus fever 174, malarial fever 8,712, smallpox 176, yellow fever 146, leprosy 81, beriberi 22, and 4,107 from phthisis pulmonalis.

*WEST INDIES—St. Christopher.*—Three weeks ended April 13, 1907. Estimated population, 49,289. Total number of deaths 82, including 5 from tuberculosis.